



Hip Evaluation Report

Report Date: 12/22/2009

Reference #: **879551**
Practice #:

Radiography Date: 3/9/2009
Date Received: 3/20/2009

Owner:
LARRY & PAULA WOODWARD
PO BOX 117
BELVUE, KS 66407-0117
UNITED STATES

PennHIP Member:
DR. WALTER RENBERG
KANSAS STATE UNIVERSITY
VETERINARY MEDICAL TEACHING HOSPITAL
1800 DENISON AVE
MANHATTAN, KS 66506-5706
UNITED STATES

ANIMAL	
AUX LAKE BRILEY (SAM) CANINE / WIREHAISED POINTING GRIFFON Date of Birth: 7/20/2006 Sex: M Weight: 65 lbs. Age: 32 mo.	Reg. #: SR36737301 Microchip: 081 865 265 AVID Tattoo:

RESULTS			
LEFT	Distraction Index (DI)	0.20	DI is less than or equal to 0.30, with no radiographic evidence of DJD.
	Degenerative Joint Disease (DJD)	None	
	Cavitation	No	
	Other Findings	Not Applicable	
RIGHT	Distraction Index (DI)	0.18	DI is less than or equal to 0.30, with no radiographic evidence of DJD.
	Degenerative Joint Disease (DJD)	None	
	Cavitation	No	
	Other Findings	Not Applicable	

Please note that the PennHIP DI is a measure of hip joint laxity, it does not allude to a "passing" or "failing" hip score.

LAXITY PROFILE RANKING

The laxity profile ranking is based on the hip with the greater laxity (DI). This interpretation is based on a cross-section of 338 CANINE animals of the WIREHAISED POINTING GRIFFON breed. The median DI for this group is 0.39.

Percentiles										
	90th	80th	70th	60th	50th	40th	30th	20th	10th	
> 90th					Median					< 10th
↑										

The chart above indicates the ranking of your animal's passive hip laxity (DI) in relation to all CANINE animals of the WIREHAISED POINTING GRIFFON breed in our database. This result means that 1) your animal's hips are tighter than approximately 100% of this group of animals (alternatively, 0% of the group has tighter hips than your animal), and 2) your animal's hip laxity is in the tighter half of the laxity profile. Breed-specific evaluations are analyzed semi-annually. Consequently, the average laxity and range of laxity for any given group will change over time.

PennHIP does not make specific breeding recommendations. Selection of sire and dam for mating is the decision of the breeder.

NOTE: As a minimum breeding criterion, we propose that breeding stock be selected from the population of animals having hip laxity in the tighter half of the breed (to the left of the median mark on the graph). Higher selection pressure equates to more rapid expected genetic change per generation.

By implementing selection based on passive hip laxity, we expect the breed average DI over the years to move toward tighter hip configuration, meaning lower hip dysplasia susceptibility. The PennHIP database permits scientific adjustment of criteria to reflect these shifts; the average laxity and range of laxity for a particular breed will change over time.